

Chapter 4

ANTHRAX

BRET K. PURCELL, PhD, MD^{*}; PATRICIA L. WORSHAM, PhD[†]; AND ARTHUR M. FRIEDLANDER, MD[‡]

INTRODUCTION AND HISTORY

THE ORGANISM

EPIDEMIOLOGY

PATHOGENESIS

CLINICAL DISEASE

Cutaneous Anthrax

Inhalational Anthrax

Oropharyngeal and Gastrointestinal Anthrax

Meningitis

DIAGNOSIS

TREATMENT

PROPHYLAXIS

Prophylactic Treatment After Exposure

Active Immunization

Side Effects

SUMMARY

^{*}Lieutenant Colonel, Medical Corps, US Army; Chief, Bacterial Therapeutics, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases, 1425 Porter Street, Fort Detrick, Maryland 21702

[†]Deputy Chief, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases, 1425 Porter Street, Fort Detrick, Maryland 21702

[‡]Colonel, Medical Corps, US Army (Ret); Senior Scientist, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases, 1425 Porter Street, Fort Detrick, Maryland 21702; and Adjunct Professor of Medicine, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Road, Bethesda, Maryland 20814

INTRODUCTION AND HISTORY

Anthrax, a zoonotic disease caused by *Bacillus anthracis*, occurs in domesticated and wild animals, primarily herbivores, including goats, sheep, cattle, horses, and swine.¹⁴ Humans usually become infected by contact with infected animals or contaminated animal products, most commonly via the cutaneous route and only rarely via the respiratory or gastrointestinal routes.^{5,6} Anthrax has a long association with human history. The fifth and sixth plagues described in Exodus may have been anthrax in domesticated animals followed by cutaneous anthrax in humans. Virgil described anthrax in domestic and wild animals in his *Georgics*, and anthrax was an economically important agricultural disease during the 16th through 18th centuries in Europe.^{7,8}

Anthrax, which is intimately associated with the origins of microbiology and immunology, was the first disease for which a microbial origin was definitively established. Robert Koch established the microbial origin for anthrax in 1876.^{9,10} Anthrax also was the first disease for which an effective live bacterial vaccine was developed; Louis Pasteur developed that vaccine in 1881.¹¹ Additionally, anthrax represents the first described occupational respiratory infectious disease. During the latter half of the 19th century, inhalational anthrax,¹² a previously unrecognized form, occurred among woolsorters in England as a result of the generation of infectious aerosols of anthrax spores under

industrial conditions from the processing of contaminated goat hair and alpaca wool.¹³

The military has long been concerned about anthrax as a potential biological weapon because anthrax spores are infectious by the aerosol route, and a high mortality rate is associated with untreated inhalational anthrax. In 1979 the largest inhalational anthrax epidemic of the 20th century occurred in Sverdlovsk, Russia. Anthrax spores were accidentally released from a military research facility located upwind from where the cases occurred. According to the accounts provided by two Soviet physicians, 96 human anthrax cases were reported, of which 79 were gastrointestinal and 17 cutaneous. The 79 gastrointestinal cases resulted in 64 deaths. Although the initial report of this event attributed the infections to a gastrointestinal source, later evidence indicated that an aerosol release of weaponized anthrax spores from a military production facility had occurred, and thus, inhalational anthrax may have been the predominant cause of these civilian casualties. Retrospective analysis using administrative name lists of compensated families, household interviews, grave markers, pathologists' notes, various hospital lists, and clinical case histories of five survivors yielded evidence of 77 anthrax cases, with 66 deaths and 11 survivors.¹⁴ Cases were also reported in animals located more than 50 km from the site.^{15,16} Polymerase chain reaction examination of tissue samples collected from 11 of the




Fig. 4-1. (a) Gram stain of a blood smear from an infected guinea pig demonstrating intracellular bacilli chains within a polymorphonuclear leukocyte. (b) Gram stain of peripheral blood smear from a nonhuman primate infected with *Bacillus anthracis*, Ames strain.

Photograph: Courtesy of Susan Welkos, PhD, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Maryland.

Photograph: Courtesy of John Ezzell, PhD, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Maryland.

victims demonstrated that virulent *B anthracis* DNA was present in all these patients, and at least five different strains of virulent anthrax were detected based on variable number tandem repeat analysis.¹⁷

Although the Sverdlovsk incident is not well known among US civilians, most people are familiar with the 2001 bioterrorist attack in the United States in letters containing dried *B anthracis* spores. The spore powder, which was sealed in letters addressed to members of the press and of Congress, was mailed through the US Postal Service.¹⁸⁻²⁰ According to the Centers for Disease Control and Prevention, 22 people contracted anthrax from the letters.^{18,21-25} Of the 11 individuals who developed inhalational anthrax, five died and six survived after intensive antimicrobial therapy. Eleven other people contracted cutaneous anthrax; all survived after treatment. Thousands of other persons received prophylaxis with antibiotics and, in some cases, postexposure vaccination.²⁶⁻²⁹ This incident profoundly affected the law enforcement, scientific, and medical communities within the United States. As a result of the attacks, there has been increased governmental and public awareness of the threat posed by *B anthracis* and other pathogens, particularly those with a potential for aerosol-mediated infection.³⁰⁻⁴³ The amount of funding budgeted to prepare and protect the nation from a bioterror attack has rapidly increased since 2001, and a significant amount of this funding has supported anthrax studies. Some of the new anthrax studies have focused on improved sample collection,




Fig. 4-2. Scanning electron micrograph of a preparation of *Bacillus anthracis* spores. Two elongated bacilli are also presented among the oval-shaped spores. Original magnification $\times 2620$.

Photograph: Courtesy of John Ezzell, PhD, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Maryland.

rapid detection/diagnosis, decontamination, and microbial forensics. Because of the ongoing terrorism threat, there has been a particular sense of urgency regarding the development and improvement of medical countermeasures, such as therapeutics, vaccines, diagnostics, and devices.

THE ORGANISM

B anthracis is a large, gram-positive, spore-forming, nonmotile bacillus ($1-1.5 \mu\text{m} \times 3-10 \mu\text{m}$) that is closely related to *B cereus* and *B thuringiensis*. The organism grows readily on sheep blood agar aerobically and is nonhemolytic under these conditions. The colonies are large, rough, and grayish white, with irregular, curving outgrowths from the margin. The organism forms a prominent capsule both in vitro in the presence of bicarbonate and carbon dioxide and in tissue in vivo. In tissue, the encapsulated bacteria occur singly or in chains of two or three bacilli (Figure 4-1). The organism does not form spores in living tissue; sporulation occurs only after the infected body has been opened

and exposed to oxygen. The spores, which cause no swelling of the bacilli, are oval and occur centrally or paracentrally (Figure 4-2). The spores are very resistant and may survive for decades in certain soil conditions. Bacterial identification is confirmed by demonstration of the protective antigen (PA) toxin component, lysis by a specific bacteriophage, detection of capsule by fluorescent antibody, and virulence for mice and guinea pigs.^{44,45} Additional confirmatory tests to identify toxin and capsule genes by polymerase chain reaction, developed as research tools, have been incorporated into the Laboratory Response Network established by the Centers for Disease Control and Prevention.⁴⁶⁻⁴⁹

EPIDEMIOLOGY

Anthrax, an organism that exists in the soil as a spore, occurs worldwide. Whether its persistence in the soil results from significant multiplication of the organism, or from cycles of bacterial amplification in infected animals whose carcasses then contaminate the

soil, remains unsettled.^{50,51} The form of the organism in infected animals is the bacillus. Sporulation occurs only when the organism in the carcass is exposed to air.

Domestic or wild animals become infected when they ingest spores while grazing on contaminated

land or eating contaminated feed. Pasteur originally reported that environmental conditions such as drought, which may promote trauma in the oral cavity on grazing, may increase the chances of acquiring anthrax.⁵² Spread from animal to animal by mechanical means—by biting flies and from one environmental site to another by nonbiting flies and by vultures—has been suggested to occur.^{51,53}

Anthrax in humans is associated with agricultural, horticultural, or industrial exposure to infected animals or contaminated animal products. In less developed countries, primarily Africa, Asia, and the Middle East, disease occurs from contact with infected domesticated animals or contaminated animal products. Contact may include handling contaminated carcasses, hides, wool, hair, and bones and ingesting contaminated meat. Cases associated with industrial exposure, rarely seen now, occur in workers processing contaminated hair, wool, hides, and bones. Direct contact with contaminated material leads to cutaneous disease, and ingestion of infected meat leads to oropharyngeal or gastrointestinal forms of anthrax. Inhalation of a sufficient quantity of spores, usually seen only during generation of aerosols in an enclosed space associated with processing contaminated wool or hair, leads to inhalational anthrax. Military research facilities have played a major role in studying and defining anthrax, as well as many other zoonotic diseases in wild and domestic animals and the subsequent infections in humans.⁵⁴

Unreliable reporting makes it difficult to estimate with accuracy the true incidence of human anthrax. It

was estimated in 1958 that between 20,000 and 100,000 cases occurred annually worldwide.⁵⁵ In more recent years, anthrax in animals has been reported in 82 countries, and human cases continue to be reported from Africa, Asia, Europe, and the Americas.⁵⁶⁻⁶⁰ In the 1996–1997 global anthrax report, there appeared to be a general decrease in anthrax cases worldwide; however, anthrax remains underdiagnosed and underreported.⁶¹ In the United States the annual incidence of human anthrax has steadily declined from about 127 cases in the early part of the 20th century to about 1 per year for the past 10 years. The vast majority of these cases have been cutaneous. Under natural conditions, inhalational anthrax is rare; before the anthrax bioterrorism event in 2001, only 18 cases had been reported in the United States in the 20th century.^{62,63} In the early part of the 20th century, inhalational anthrax cases were reported in rural villagers in Russia who worked with contaminated sheep wool inside their homes.⁶⁴ However, in recent years a significant decrease occurred in anthrax cases in domestic animals in east Russia. Five inhalational anthrax cases occurred in woolen mill workers in New Hampshire in the 1950s.⁶⁵ During economic hardship and disruption of veterinary and human public health practices (eg, during wartime), large anthrax epidemics have occurred. The largest reported human anthrax epidemic occurred in Zimbabwe from 1978 through 1980, with an estimated 10,000 cases. Essentially all cases were cutaneous, including rare gastrointestinal disease cases and eight inhalational anthrax cases, although no autopsy confirmation was reported.⁶⁶

PATHOGENESIS

B anthracis possesses two protein exotoxins, known as the lethal toxin and the edema toxin; an antiphagocytic capsule; and other known and putative virulence factors.⁶⁷ The role of the capsule in pathogenesis was demonstrated in the early 1900s, when anthrax strains lacking a capsule were shown to be avirulent.⁶⁸ In more recent years, the genes encoding synthesis of the capsule were encoded on the 96-kilobase (kb) plasmid known as pXO2. Molecular analysis revealed that strains cured of this plasmid no longer produced the capsule and were attenuated, thus confirming the critical role of the capsule in virulence.⁶⁹ The capsule is composed of a polymer of poly-D-glutamic acid, which confers resistance to phagocytosis and may contribute to the resistance of anthrax to lysis by serum cationic proteins.⁷⁰ Capsule production is necessary for dissemination to the spleen in a murine inhalational anthrax model.⁷¹ Recently, the capsule has also been the focus of several efforts to develop new generation

anthrax vaccines.⁷²⁻⁷⁴ Evidence indicates that the capsule may enhance the protection afforded by PA-based vaccines against anthrax if opsonizing antibodies are produced.⁷⁴

Koch first suggested the importance of toxins in his initial studies on anthrax. In 1954 Smith and Keppie⁷⁵ demonstrated a toxic factor in the serum of infected animals that was lethal when injected into other animals. The role of toxins in virulence and immunity was firmly established by many researchers in the ensuing years.⁷⁶⁻⁷⁸ Advances in molecular biology in the past decade have produced a more complete understanding of the biochemical mechanisms of action of the toxins and have begun to provide a more definitive picture of their role in the pathogenesis of the disease.

Two protein exotoxins, known as the lethal toxin and edema toxin, are encoded on a 182-kb plasmid (pXO1), distinct from that coding for the capsule. In an

environment of increased bicarbonate, carbon dioxide, and increased temperature, such as is found in the infected host, transcription of the genes encoding these and other virulence-associated gene products is enhanced.^{67,79-82} A complex regulatory cascade controlled in large part by the *atxA* and *acpA* genes encoded on the toxin plasmid pXO1 and pXO2, respectively, directs the production of virulence factors in response to these environmental signals.^{83,84}

Recently, a retrospective study identified an isolate of *B cereus* that carried a plasmid homologous to the anthrax toxin plasmid pXO1. This strain was obtained from a patient with symptoms similar to inhalational anthrax.⁸⁵ This finding led to considerable concern because “anthrax toxin” sequences are considered unique to *B anthracis*. Although a polyglutamate capsule was not produced, sequences encoding a polysaccharide capsule were present on a smaller plasmid. The possibility of false positives from toxin-based identification tests should be considered because many diagnostic schemes have focused on toxin genes and gene products. The virulence of this isolate has not yet been extensively studied, and the role of the lethal and edema toxins in the pathogenesis of this strain is unknown. Likewise, the incidence of such strains in nature is unclear. Because *B cereus* is hemolytic and resistant to the anthrax-specific gamma bacteriophage, such isolates would not typically be tested for the presence of genes encoding anthrax toxin, especially because *B cereus* is often regarded as an environmental contaminant.⁸⁵ Other human cases of anthrax-like *B cereus* infections have been reported.^{86,87}

The anthrax toxins, like many bacterial and plant toxins, possess two components: (1) a cell-binding, pore-forming, or B, domain; and (2) an active, or A, domain that has the toxic and, usually, the enzymatic activity (Figure 4-3). The B and A anthrax toxin components are synthesized from different genes and are secreted as noncovalently linked proteins. The anthrax toxins are unusual in that the B protein, PA, is shared by both toxins. Thus, the lethal toxin is composed of the PA₆₃ (MW 63,000 after cleavage from a MW 83,000 protein) heptamer combined with a second protein, which is known as the lethal factor (LF [MW 90,000]), and the edema toxin is composed of PA complexed with the edema factor (EF [MW 89,000]). Each of the three toxin proteins—the B protein and both A proteins—individually is without biological activity. The critical role of the toxins in pathogenesis was established when it was shown that deletion of the toxin-encoding plasmid pXO1^{69,88} or the PA gene alone⁸⁹ attenuates the organism. The lethal toxin appears to be more important for virulence in a mouse model than the edema toxin.⁹⁰ Crude toxin

preparations have been shown to impair neutrophil chemotaxis⁹¹ and phagocytosis.⁷⁰

The edema toxin causes edema when injected into the skin of experimental animals and is likely responsible for the marked edema often present at bacterial replication sites.^{92,93} This toxin is a calmodulin-dependent adenylate cyclase that impairs phagocytosis and priming for the respiratory burst in neutrophils; it also inhibits the production of interleukin-6 and tumor necrosis factor by monocytes, which may further weaken host resistance.⁹⁴⁻⁹⁶ Edema toxin also impairs dendritic cell function and appears to act with lethal toxin to suppress the innate immune response.⁹⁷

The lethal toxin is a zinc metalloprotease that is lethal for experimental animals^{92,93,98} and is directly cytolytic for macrophages, causing release of the potentially toxic cytokines interleukin-1 and tumor




Fig. 4-3. Composition of anthrax lethal protein toxin. Molecular models of the protective antigen (PA)₆₃ heptamer and the PA₆₃ heptamer-lethal factor (LF) complex. (a, b) Side and top views of PA₆₃ heptamer (green) bound to three LF molecules (yellow). (c, d) The surface renderings are colored according to the negative (red) and positive (blue) electrostatic surface potential. (c) Top view of the PA₆₃ heptamer. The yellow box highlights the protomer-protomer interface and where LF binds to heptameric PA. (d) A hypothetical PA₆₃ heptamer-LF interface.

Photographs: Courtesy of Kelly Halverson, PhD, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Maryland.

necrosis factor.⁹⁹ In in-vitro models, lethal toxin cleaves members of the mitogen-activated protein kinase (MAPK) kinase family, which are an integral part of a phosphorelay system that links surface receptors to transcription of specific genes within the nucleus. Thus, lethal toxin interferes with the MAPK signaling pathways necessary for a multitude of normal cell functions.¹⁰⁰ In macrophage and dendritic cell models, lethal toxin leads to inhibition of proinflammatory cytokines, downregulation of costimulatory molecules, and ineffective T-cell priming.¹⁰⁰⁻¹⁰³ It also appears to promote apoptosis of endothelial cells lining the vascular system in vitro, leading to speculation that lethal toxin-induced barrier dysfunction leads to the vascular permeability changes accompanying systemic anthrax infection.¹⁰⁴ Effects on hormone receptors, including glucocorticoids, have also been reported. Although much of the information regarding lethal toxin activity has been obtained from animal-derived cell culture models, Fang et al recently reported that, in vitro, lethal toxin inhibits MAPK kinase dependent interleukin-2 production and proliferative responses in human CD4⁺ T cells.¹⁰⁵ The in-vivo targets for these toxins await confirmation; however, both lethal and edema toxins contribute significantly to suppression of the innate immune system.

Recent studies in cell culture models have given a clearer understanding of the molecular interactions of the toxin proteins.¹⁰⁰ PA first binds, most likely by a domain at its carboxy-terminus, to a specific cell receptor.¹⁰⁶⁻¹⁰⁸ Two proteins have been proposed as the PA receptor: (1) anthrax toxin receptor protein, ANTX1; and (2) capillary morphogenesis protein, CMG2.^{109,110} Although their natural ligands have not been identified, both receptors have a von Willibrand factor type A domain that appears to interact with PA. Once bound, PA is cleaved by a furin-like protease, resulting in retention of a 63-kilodalton (kd) fragment of PA on the cell surface.^{111,112} This cleavage promotes formation of PA heptamers and creates a binding site on PA to which up to three molecules of the LF and the EF can bind with high affinity.^{101,113} Heptamerization stimulates endocytosis of PA (or PA-EF/LF complexes), which is then delivered to early endosomes.¹¹⁴ The mildly acidic pH of the endosome triggers membrane insertion of the heptameric PA into intraluminal vesicles.¹¹⁵ EF

and LF are translocated into the lumen of the vesicle and are thereby protected from lysosomal proteases.¹¹⁵ The toxins are then translocated via endosomal carrier vesicles to the cell cytosol, where they express their toxic activity.¹¹⁵

The processes leading to toxin activity in the infected animal may be more complicated because the toxin proteins appear to exist in the serum as a complex of PA and EF/LF.¹¹⁶ The proteolytic activation of PA necessary to form lethal or edema toxin may occur in interstitial fluid or serum rather than on the cell surface.¹¹⁶ The lethal or edema toxin may then bind to target cells and be internalized. This theory was recently bolstered by Panchal et al who demonstrated that purified LF complexed with the PA(110) heptamer cleaved both a synthetic peptide substrate and endogenous MAPK kinase substrates and killed susceptible macrophage cells.¹¹⁷ In addition, complexes of the heptameric PA(110)-LF found in the plasma of infected animals showed functional activity.¹¹⁷ Terminally, toxin is present in high concentrations in the blood, but the molecular mechanisms that cause death remain unknown.⁷⁶

Infection begins when the spores are inoculated through the skin or mucosa. Spores are ingested at the local site by macrophages, in which they germinate to the vegetative bacillus with production of capsule and toxins. At these sites, the bacteria proliferate and produce the edema and lethal toxins that impair host leukocyte function and lead to the distinctive pathological findings: edema, hemorrhage, tissue necrosis, and a relative lack of leukocytes. In inhalational anthrax, the spores are ingested by alveolar macrophages, which transport them to the regional tracheobronchial lymph nodes, where germination occurs.¹¹⁸ Once in the tracheobronchial lymph nodes, the local production of toxins by extracellular bacilli generates the characteristic pathology picture: massive hemorrhagic, edematous, and necrotizing lymphadenitis; and mediastinitis (the latter is almost pathognomonic of this disease).¹¹⁹ The bacilli can then spread to the blood, leading to septicemia with seeding of other organs and frequently causing hemorrhagic meningitis. Death is the result of respiratory failure associated with pulmonary edema, overwhelming bacteremia, and often meningitis.

CLINICAL DISEASE

The military seeks to defend against anthrax used as an inhalational biological weapon. However, other anthrax forms are more likely to be seen by medical officers—particularly when deployed to third world countries—and are therefore included for completeness.

Cutaneous Anthrax

More than 95% of anthrax cases are cutaneous.^{120,121} After inoculation, the incubation period is 1 to 5 days. The disease first appears as a small papule

that progresses over a day or two to a vesicle containing serosanguineous fluid with many organisms and a paucity of leukocytes. Histopathology findings consist of varying degrees of ulceration, vasculitis, perivascular inflammation, coagulative necrosis, hemorrhage, and edema.¹²² The vesicle, which may be 1 to 2 cm in diameter, ruptures, leaving a necrotic ulcer (Figure 4-4). Satellite vesicles may also be present. The lesion is usually painless, and varying degrees of edema may be present around it.¹²³ The edema may occasionally be massive, encompassing the entire face or limb, which is described by the term “malignant edema.” Patients usually have fever, malaise, and headache, which may be severe in those with extensive edema. There may also be local lymphadenitis. The ulcer base develops a characteristic black eschar, and after 2 to 3 weeks the eschar separates, often leaving a scar and sometimes requiring surgical reconstruction.^{124,125} Septicemia is rare, and with treatment mortality should be less than 1%.^{124,126-128} In addition, no age-related risk factor appears to be associated with cutaneous human anthrax.¹²⁹

Inhalational Anthrax

Inhalational anthrax begins after an incubation period of 1 to 6 days with nonspecific symptoms of malaise, fatigue, myalgia, and fever.¹³⁰⁻¹³³ A nonproductive cough and mild chest discomfort may also occur. These symptoms usually persist for 2 or 3 days, and in some cases there may be a short period of improvement. Then a sudden onset of increasing respiratory distress with dyspnea, stridor, cyanosis, increased chest pain, and diaphoresis occurs. Associated edema of the chest and neck may also be present. Chest radiograph examination usually shows the characteristic widening of the mediastinum from necrosis and hemorrhage of the lymph nodes and surrounding tissues, often with associated pleural effusions (Figure 4-5). In the 2001 bioterrorist event, the pleural effusions were initially small but rapidly progressed and persisted despite effective antibiotic therapy.^{134,135} The effusions were predominantly serosanguineous and immunohistochemistry revealed the presence of *B anthracis* cell walls and capsule antigens. Effusion fluid from deceased patients who had received fewer than 55 hours of




Fig. 4-4. Cutaneous lesions of anthrax. (a) Ulcer with vesicle ring. (b) Black eschar with surrounding erythema. (c) Marked edema of extremity secondary to anthrax edema toxin with multiple black eschar.

Photographs: Courtesy of the Centers for Disease Control and Prevention, Atlanta, Georgia. www.bt.cdc.gov/agent/anthrax/anthrax-images/cutaneous.asp.




Fig. 4-5. (a) Frontal chest radiograph reveals mediastinal and hilar widening, bilateral pleural effusions, and decreased lung volumes. (b) Chest axial computed tomography (CT) (mediastinal window) shows enlarged, hyperdense subcarinal (arrow) and left hilar (arrowhead) lymph nodes, compatible with intranodal hemorrhage. (c) On lung window CT, peribronchovascular consolidation (curved arrow) reflects lymphatic spread of anthrax infection.

Radiologic Images: Courtesy of JR Galvin, MD and AA Frazier, MD, Department of Radiologic Pathology, Armed Forces Institute of Pathology, Washington, DC.




Fig. 4-6. Meningitis with subarachnoid hemorrhage in a man from Thailand who died 5 days after eating undercooked carabao (water buffalo).

Reproduced from: Binford CH, Connor DH, eds. *Pathology of Tropical and Extraordinary Diseases*. Vol 1. Washington, DC: Armed Forces Institute of Pathology; 1976: 121. AFIP Negative 75-12374-3.

antibiotic therapy revealed the presence of bacilli.¹³⁶ Polymerase chain reaction analysis of the pleura fluid was also positive for *B anthracis* DNA.¹³⁷ Pneumonia has not been a consistent finding but can occur in some patients⁵ and may be attributed to intravascular edema and hyaline membrane formation.¹³⁶ Although inhalational anthrax cases have been rare in this century, except for the 11 cases arising from the anthrax letters in 2001, several cases have occurred in patients with underlying pulmonary disease, suggesting that this condition may increase susceptibility to the disease.⁵² Meningitis is present in up to 50% of cases, and some patients may present with seizures. The onset of respiratory distress is followed by the rapid onset of shock and death within 24 to 36 hours. Mortality had been essentially 100% in the absence of appropriate treatment; however, during 2001 the mortality rate was 45%.^{134,135}

Oropharyngeal and Gastrointestinal Anthrax

Oropharyngeal and gastrointestinal anthrax result from ingesting infected meat that has not been sufficiently cooked.¹³⁸ After an incubation period of 2 to

5 days, patients with oropharyngeal disease present with severe sore throat or a local oral or tonsillar ulcer, usually associated with fever, toxicity, and swelling of the neck resulting from cervical or submandibular lymphadenitis and edema. Dysphagia and respiratory distress may also be present. Gastrointestinal anthrax begins with nonspecific symptoms of nausea, vomiting, and fever; in most cases severe abdominal pain follows. The presenting sign may be an acute abdomen, which may be associated with hematemesis, massive ascites, and bloody diarrhea. Mortality in

both forms may be as high as 50%, especially in the gastrointestinal form.

Meningitis

Meningitis may occur after bacteremia as a complication of any of the other clinical forms of the disease.¹³⁹ Meningitis may also occur—rarely—without a clinically apparent primary focus, and it is often hemorrhagic, which is important diagnostically, and almost always fatal (Figure 4-6).

DIAGNOSIS

The most critical aspect in making an anthrax diagnosis is a high index of suspicion associated with a compatible history of exposure. Cutaneous anthrax should be considered after a painless pruritic papule, vesicle, or ulcer develops—often with surrounding edema—and then becomes a black eschar. With extensive or massive edema, such a lesion is almost pathognomonic. Gram stain or culture of the lesion usually confirms the diagnosis. Bacterial culture tests include colony morphology on sheep blood agar plates incubated at 35°C to 37°C for 15 to 24 hours. *B anthracis* colonies are 2 to 5 mm in diameter, flat or slightly convex, irregularly round with possible comma-shaped (“Medusa-head”) projections with a ground-glass appearance (Figure 4-7). The colonies

tend to have tenacious consistency when moved with a bacterial loop and are not β -hemolytic. The bacteria appear as gram-positive, 1 to 8 μm long and 1 to 1.5 μm wide bacilli. India ink staining reveals capsulated bacteria. A motility test should be performed either by wet mount or motility media; *B anthracis* is nonmotile. Gamma bacteriophage lysis and direct fluorescent antibody tests are performed at Level D laboratories as confirmatory tests (see Figure 4-7 and Figure 4-8). Commercial polymerase chain reaction kits specific for the *B anthracis* pX01 and pX02 plasmids are also available to assist in identification of this organism. The differential diagnosis should include tularemia, staphylococcal or streptococcal disease, and orf (a viral disease of sheep and goats transmissible to humans).




Fig. 4-7. (a) Isolated colonies of *Bacillus anthracis* on sheep blood agar plate. (b) Detection of *B anthracis* using specific gamma-phage mediated cell-lysis.

Photographs: Courtesy of Bret K Purcell, PhD, MD, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases and the Defense Threat Reduction Agency/Threat Agent Detection and Response Program, National Center for Disease Control, Tbilisi, Georgia, 2005.




Fig. 4-8. Direct fluorescent antibody stain of *Bacillus anthracis* capsule.

Photograph: Courtesy of David Heath, PhD, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases and the Defense Threat Reduction Agency/Threat Agent Detection and Response Program, National Center for Disease Control, Tbilisi, Georgia, 2005.

The diagnosis of inhalational anthrax is difficult, but the disease should be suspected with a history of exposure to a *B anthracis*-containing aerosol. The early symptoms are nonspecific¹³¹⁻¹³³ and include fever, chills, dyspnea, cough, headache, vomiting, weakness, myalgias, abdominal pain, and chest or pleuritic pain. This stage of the disease may last from hours to a few days. However, the development of respiratory distress in association with radiographic evidence of a widened mediastinum resulting from hemorrhagic mediastinitis and the presence of hemorrhagic pleural effusion or hemorrhagic meningitis should suggest the diagnosis. Contrast-enhanced computer tomography images reveal diffuse hemorrhagic mediastinal and

hilar adenopathy with edema, perihilar infiltrates, bronchial mucosal thickening, hemorrhagic pleural, and pericardial effusions.¹⁴⁰ During the later stages of the disease patients develop sudden fever, dyspnea, diaphoresis, cyanosis, hypotension, shock, and death.¹³¹ Blood culture should demonstrate growth in 6 to 24 hours if the patient has not received antibiotics before collection, and Gram stain of peripheral blood smears often reveals large bacilli in later stages of disease. Sputum examination is not helpful in making the diagnosis because pneumonia is usually not a feature of inhalational anthrax.

Gastrointestinal anthrax is difficult to diagnose because of its rarity and nonspecific symptoms including nausea, vomiting, anorexia, and fever. As the disease progresses, patients often develop acute, severe abdominal pain; hematemesis; and bloody diarrhea. Diagnosis is usually considered only with a history of ingesting contaminated meat in the setting of an outbreak. Microbiological cultures do not help confirm the diagnosis. The diagnosis of oropharyngeal anthrax can be made from the clinical and physical findings in a patient with the appropriate epidemiological history. Sore throat, dysphagia, hoarseness, cervical lymphadenopathy, and edema as well as fever are often presenting symptoms.^{133,141,142}

Meningitis resulting from anthrax is clinically indistinguishable from meningitis attributable to other etiologies. An important distinguishing feature is that the cerebral spinal fluid is hemorrhagic in as many as 50% of cases. The diagnosis can be confirmed by identifying the organism in cerebral spinal fluid by microscopy, culture, or both.

Serology is generally only useful in making a retrospective diagnosis. Antibody to PA or the capsule develops in 68% to 93%¹⁴³⁻¹⁴⁶ of reported cutaneous anthrax cases and 67% to 94%^{145,146} of reported oropharyngeal anthrax cases. A positive skin test to anthraxin (an undefined antigen derived from acid hydrolysis of the bacillus that was developed and evaluated in the former Soviet Union) has also been reported¹⁴⁷ to help with the retrospective diagnosis of anthrax. Western countries have limited experience with this test.¹⁴⁸

TREATMENT

Cutaneous anthrax without toxicity or systemic symptoms may be treated with oral penicillin if the infection did not originate with a potential aerosol exposure. However, if an inhalational exposure is also suspected, ciprofloxacin or doxycycline is recommended as first-line therapy.^{131,149} Effective therapy reduces edema and systemic symptoms but does not change the evolution of the skin lesion. Treatment should be continued for 7 to 10 days, unless inhalational exposure

is suspected; then treatment should be continued for 60 days. However, recent studies of the 2001 bioterrorism event have identified problems associated with prolonged treatment, mass prophylaxis, and medication compliance.¹⁵⁰⁻¹⁵⁴ Amoxicillin is recommended for patients who cannot take fluoroquinolones or tetracycline-class drugs; however, increasing evidence shows that *B anthracis* possesses β -lactamase genes that may reduce the efficacy of this treatment.¹⁵⁵⁻¹⁶⁰ In addition, if a bioter-

rorism event occurs, the bacterial strains used may be intentionally antibiotic resistant or genetically modified to confer resistance to one or more antibiotics.

Tetracycline, erythromycin, and chloramphenicol have also been used successfully¹⁶¹ for treating rare cases caused by naturally occurring penicillin-resistant organisms. Additional antibiotics shown to be active in vitro include gentamicin, cefazolin, cephalothin, vancomycin, clindamycin, and imipenem.¹⁶²⁻¹⁶⁵ These drugs should be effective in vivo, but there is no reported clinical experience. Experimental infections using the inhalational mouse model have demonstrated significant efficacy using these additional antibiotics.

Inhalational, oropharyngeal, and gastrointestinal anthrax should be treated with intravenous therapy using two or more antibiotics. The therapy should initially include a fluoroquinolone or doxycycline with one or more of the following antibiotics: clindamycin, rifampin, penicillin, ampicillin, vancomycin, aminoglycosides, chloramphenicol, imipenem, clarithromycin, and linezolid.^{131,149} Patients often require intensive care unit support, including appropriate vasopressors, oxygen, and other supportive therapy, because of the disease's severity and rapid onset. Recommendations for treatment during pregnancy and for pediatric populations follow similar guidelines.^{149,159}

PROPHYLAXIS

Prophylactic Treatment After Exposure

Experimental evidence¹⁶⁶ has demonstrated that treatment with antibiotics (including ciprofloxacin, doxycycline, and penicillin) beginning 1 day after exposure to a lethal aerosol challenge with anthrax spores can significantly protect against death. Combining antibiotics with active vaccination provides the optimal protection. Recent analysis has suggested postexposure vaccination may shorten the duration of antibiotic prophylaxis, providing the least expensive and most effective strategy to counter a bioterrorism event.¹⁶⁷⁻¹⁶⁹

Active Immunization

BioPort Corporation (Lansing, Michigan) produces the only licensed human vaccine against anthrax, Anthrax Vaccine Adsorbed (BioThrax). This vaccine is made from sterile filtrates of microaerophilic cultures of an attenuated, unencapsulated, nonproteolytic strain (V770-NP1-R) of *B anthracis*. The filtrate, containing predominantly 83-kDa PA, is adsorbed to 1.2 mg/mL of aluminum hydroxide in 0.85% sodium chloride. The final product also contains 100 µg/mL of formaldehyde and 25 µg/mL of benzethonium chloride as preservatives. Some vaccine lots contain small amounts of LF and lesser amounts of EF, as determined by antibody responses in vaccinated animals,^{64,170,171} although this antibody response has not been reported in the limited observations in human vaccinees.¹⁷² Although PA is an effective immunogen,¹⁷³ it is unknown whether the small amounts of LF or EF in some lots of the vaccine contribute to its protective efficacy. The potency of vaccine lots is determined by showing protection of parenterally challenged guinea pigs. An in-vitro assay for vaccine potency is being developed.¹⁷⁴ There is no characterization of the amount and form of the PA or other toxin components in the vaccine. The vaccine

is stored at 2°C to 8°C. The recommended schedule for vaccination is 0.5 mL given subcutaneously over the deltoid muscle at 0, 2, and 4 weeks, followed by boosters of 0.5 mL at 6, 12, and 18 months. Annual boosters are recommended if the potential for exposure continues.

The vaccine should be given to industrial workers exposed to potentially contaminated animal products imported from countries in which animal anthrax remains uncontrolled. These products include wool, goat hair, hides, and bones. People in direct contact with potentially infected animals and laboratory workers should also be vaccinated. Vaccination is also indicated for protection against anthrax use in biological warfare. Recommendations have been made for anthrax vaccine use in the United States.^{175,176} More than 500,000 US military personnel have received the licensed anthrax vaccine adsorbed (AVA) vaccine, and no unusual rates of serious adverse events have been noted.¹⁷⁷ Additional studies also support the safety of the anthrax vaccine.¹⁷⁸⁻¹⁸⁶ The next generation vaccine, recombinant PA, may afford equivalent protection with a decrease in reactogenicity.

A live attenuated, unencapsulated spore vaccine is used for humans in the former Soviet Union. The vaccine is given by scarification or subcutaneously. Its developers claim that it is reasonably well tolerated and shows some degree of protective efficacy against cutaneous anthrax in clinical field trials.¹⁴⁷ New attenuated vaccines developed in the United States are being evaluated for efficacy in inhalational anthrax animal models.¹⁸⁷

In the United States vaccination with the licensed vaccine induced an immune response, measured by indirect hemagglutination, to PA in 83% of vaccinees 2 weeks after the first three doses,¹⁸⁸ and in 91% of those tested after receiving two or more doses.¹⁴⁴ One hundred percent of the vaccinees developed a rise in titer in response to the yearly booster dose. When tested by

an enzyme-linked immunosorbent assay (ELISA), the current serologic test of choice, more than 95% of vaccinees seroconvert after the initial three doses.^{172,189}

A rough correlation exists between antibody titer to PA and protection of experimental animals from infection after vaccination with the human vaccine. However, the exact relationship between antibody to PA as measured in these assays and immunity to infection remains obscure because the live attenuated Sterne veterinary vaccine (made from an unencapsulated, toxin-producing strain) protects animals better than the human vaccine, yet it induces lower levels of antibody to PA.¹⁷⁰⁻¹⁷²

The protective efficacy of experimental PA-based vaccines produced from sterile culture filtrates of *B anthracis* was clearly demonstrated by various animal models and routes of challenge.^{67,190} A placebo-controlled clinical trial was conducted with a vaccine similar to the currently licensed US vaccine.¹⁹¹ This field-tested vaccine was composed of the sterile, cell-free culture supernatant from an attenuated, unencapsulated strain of *B anthracis*, different from that used to produce the licensed vaccine and grown under aerobic, rather than microaerophilic, conditions.¹⁹² This vaccine was precipitated with alum rather than adsorbed to aluminum hydroxide. The study population worked in four mills in the northeastern United States where *B anthracis*-contaminated imported goat hair was used. The vaccinated group, compared to a placebo-inoculated control group, was afforded 92.5% protection against cutaneous anthrax, with a lower 95% confidence limit of 65% effectiveness. There were insufficient inhalational anthrax cases to determine whether the vaccine was effective. This same vaccine was previously shown to protect rhesus monkeys and other animal models against an aerosol exposure

to anthrax spores.¹⁹²⁻¹⁹⁸ No controlled clinical trials in humans of the efficacy of the currently licensed US vaccine have been conducted. This vaccine has been extensively tested in animals and has protected guinea pigs against both an intramuscular^{171,172,195} and an aerosol challenge.¹⁷⁰ The licensed vaccine has also been shown to protect rhesus monkeys against an aerosol challenge.^{166,195,196,198}

Side Effects

In two different studies, the incidence of significant local and systemic reactions to the vaccine used in the placebo-controlled field trial was 2.4% to 2.8%⁶⁶ and 0.2% to 1.3%.¹⁹² The vaccine licensed in the United States is reported to have a similar incidence of reactions.^{189,199} Local reactions considered significant include induration, erythema in an area larger than 5 cm in diameter, edema, pruritus, warmth, and tenderness. These reactions peak at 1 to 2 days and usually resolve within 2 to 3 days after they peak. Rare reactions include edema extending from the local site to the elbow or forearm, and a small, painless nodule that may persist for weeks. A recent study has indicated that frequency of local reactions could be significantly reduced by administering the vaccine over the deltoid muscle instead of the triceps.¹⁷⁷ People who have recovered from a cutaneous infection with anthrax may have severe local reactions from being vaccinated.¹⁹¹ Systemic reactions are characterized by flu-like symptoms, mild myalgia, arthralgia, headache, and mild-to-moderate malaise that last for 1 to 2 days.

There are no long-term sequelae of local or systemic reactions and no suggestion of a high frequency or unusual pattern of serious adverse events.^{177,182,183,200}

SUMMARY

Anthrax is a zoonotic disease that occurs in domesticated and wild animals. Humans become infected by contact with infected animals or contaminated products. Under natural circumstances, infection occurs by the cutaneous route and only rarely by the inhalational or gastrointestinal routes.

An aerosol exposure to spores causes inhalational anthrax, which is of military concern because of its potential for use as a biological warfare agent. Aerosol exposure begins with nonspecific symptoms followed in 2 to 3 days by the sudden onset of respiratory distress

with dyspnea, cyanosis, and stridor; it is rapidly fatal. Radiography of the chest often reveals characteristic mediastinal widening, indicating hemorrhagic mediastinitis. Hemorrhagic meningitis frequently coexists. Given the rarity of the disease and its rapid progression, it is difficult to diagnose inhalational anthrax. Treatment consists of massive doses of antibiotics and supportive care. Postexposure antibiotic prophylaxis is effective in laboratory animals and should be instituted as soon as possible after exposure. A licensed, antigen-based, nonviable vaccine is available for human use.

REFERENCES

1. Hugh-Jones ME, de Vos V. Anthrax and wildlife. *Rev Sci Tech.* 2002;21:359–383.
2. Bales ME, Dannenberg AL, Brachman PS, Kaufmann AF, Klatsky PC, Ashford DA. Epidemiologic response to anthrax outbreaks: field investigations, 1950–2001. *Emerg Infect Dis.* 2002;8:1163–1174.
3. Woods CW, Ospanov K, Myrzabekov A, Favorov M, Plikaytis B, Ashford DA. Risk factors for human anthrax among contacts of anthrax-infected livestock in Kazakhstan. *Am J Trop Med Hyg.* 2004;71:48–52.
4. Dragon DC, Bader DE, Mitchell J, Wollen N. Natural dissemination of *Bacillus anthracis* spores in northern Canada. *Appl Environ Microbiol.* 2005;71:1610–1615.
5. Stone SE. Cases of malignant pustule. *Boston Med Surg J.* 1868;1:19–21.
6. Macher A. Industry-related outbreak of human anthrax, Massachusetts, 1868. *Emerg Infect Dis.* 2002;8:1182.
7. Dirckx JH. Virgil on anthrax. *Am J Dermatopathol.* 1981;3:191–195.
8. Morens DM. Characterizing a “new” disease: epizootic and epidemic anthrax, 1769–1780. *Am J Public Health.* 2003;93:886–893.
9. Koch R. Die Aetiologie der Milzbrand-Krankheit, begründet auf die Entwicklungsgeschichte des *Bacillus anthracis*. *Beiträge zur Biologie der Pflanzen.* 1876;2:277–310.
10. Jay V. The legacy of Robert Koch. *Arch Pathol Lab Med.* 2001;125:1148–1149.
11. Pasteur L, Chamberland R, Roux E. Compte rendu sommaire des expériences faites à Pouilly-le-Fort, près Melun, sur la vaccination charbonneuse. *Comptes Rendus des séances De L’Académie des Sciences.* 1881;92:1378–1383.
12. LaForce FM. Woolsorters’ disease in England. *Bull N Y Acad Med.* 1978;54:956–963.
13. Carter T. The dissemination of anthrax from imported wool: Kidderminster 1900–14. *Occup Environ Med.* 2004;61:103–107.
14. Meselson M, Guillemin J, Hugh-Jones M, et al. The Sverdlovsk anthrax outbreak of 1979. *Science.* 1994;266:1202–1208.
15. Abramova FA, Grinberg LM, Yampolskaya OV, Walker DH. Pathology of inhalational anthrax in 42 cases from the Sverdlovsk outbreak of 1979. *Proc Natl Acad Sci U S A.* 1993;90:2291–2294.
16. Walker DH, Yampolska L, Grinberg LM. Death at Sverdlovsk: what have we learned? *Am J Pathol.* 1994;144:1135–1141.
17. Jackson PJ, Hugh-Jones ME, Adair DM, et al. PCR analysis of the tissue samples from the 1979 Sverdlovsk anthrax victims: the presence of multiple *Bacillus anthracis* strains in different victims. *Proc Natl Acad Sci U S A.* 1998;95:1224–1229.
18. Jernigan DB, Raghunathan PL, Bell BP, et al. Investigation of bioterrorism-related anthrax, United States, 2001: epidemiologic findings. *Emerg Infect Dis.* 2002;8:1019–1028.
19. Greene CM, Reefhuis J, Tan C, et al. Epidemiologic investigations of bioterrorism-related anthrax, New Jersey, 2001. *Emerg Infect Dis.* 2002;8:1048–1055.
20. Hsu VP, Lukacs SL, Handzel T, et al. Opening a *Bacillus anthracis*-containing envelope, Capitol Hill, Washington, DC: the public health response. *Emerg Infect Dis.* 2002;8:1039–1043.
21. Traeger MS, Wiersma ST, Rosenstein NE, et al. First case of bioterrorism-related inhalational anthrax in the United States, Palm Beach County, Florida, 2001. *Emerg Infect Dis.* 2002;8:1029–1034.

22. Maillard JM, Fischer M, McKee KT, Turner LF, Cline JS. First case of bioterrorism-related inhalational anthrax, Florida, 2001: North Carolina investigation. *Emerg Infect Dis.* 2002;8:1035–1038.
23. Holtz TH, Ackelberg JA, Kool JL, et al. Isolated case of bioterrorism-related inhalational anthrax, New York City, 2001. *Emerg Infect Dis.* 2003;9:689–696.
24. Griffith KS, Mead P, Armstrong GL, et al. Bioterrorism-related inhalational anthrax in an elderly woman, Connecticut, 2001. *Emerg Infect Dis.* 2003;9:681–688.
25. Dewan PK, Fry AM, Laseron K, et al. Inhalational anthrax outbreak among postal workers, Washington, DC, 2001. *Emerg Infect Dis.* 2002;8:1066–1072.
26. Guarner J, Jernigan JA, Shieh WJ, et al. Pathology and pathogenesis of bioterrorism-related inhalational anthrax. *Am J Pathol.* 2003;163:701–709.
27. Shieh WJ, Guarner J, Paddock C, et al. The critical role of pathology in the investigation of bioterrorism-related cutaneous anthrax. *Am J Pathol.* 2003;163:1901–1910.
28. Tierney BC, Martin SW, Franzke LH, et al. Serious adverse events among participants in the Centers for Disease Control and Prevention's Anthrax Vaccine and Antimicrobial Availability Program for persons at risk for bioterrorism-related inhalational anthrax. *Clin Infect Dis.* 2003;37:905–911.
29. Quinn CP, Dull PM, Semenova V, et al. Immune responses to *Bacillus anthracis* protective antigen in patients with bioterrorism-related cutaneous or inhalation anthrax. *J Infect Dis.* 2004;190:1228–1236.
30. Brachman PS. Bioterrorism: an update with a focus on anthrax. *Am J Epidemiol.* 2002;155:981–987.
31. Utrup LJ, Frey AH. Fate of bioterrorism-relevant viruses and bacteria, including spores, aerosolized into an indoor air environment. *Exp Biol Med.* 2004;229:345–350.
32. Fennelly KP, Davidow AL, Miller SL, Connell N, Ellner JJ. Airborne infection with *Bacillus anthracis*—from mills to mail. *Emerg Infect Dis.* 2004;10:996–1002.
33. Webb GF, Blaser M. Mailborne transmission of anthrax: modeling and implications. *Proc Natl Acad Sci U S A.* 2002;99:7027–7032.
34. Weiss CP, Intrepido AJ, Miller AK, et al. Secondary aerosolization of viable *Bacillus anthracis* spores in a contaminated US Senate office. *JAMA.* 2002;288:2853–2858.
35. Sanderson WT, Stoddard RR, Echt AS, et al. *Bacillus anthracis* contamination and inhalational anthrax in a mail processing and distribution center. *J Appl Microbiol.* 2004;96:1048–1056.
36. Darling RG, Catlett CL, Huebner KD, Jarrett DG. Threats in bioterrorism. I: CDC category A agents. *Emerg Med Clin North Am.* 2002;20:273–309.
37. Nicas M, Hubbard A. A risk analysis approach to selecting respiratory protection against airborne pathogens used for bioterrorism. *AIHA J.* 2003;64:95–101.
38. Webb GF. A silent bomb: the risk of anthrax as a weapon of mass destruction. *Proc Natl Acad Sci U S A.* 2003;100:4355–4356.
39. Reshetin VP, Regens JL. Simulation modeling of anthrax spore dispersion in a bioterrorism incident. *Risk Anal.* 2003;23:1135–1145.
40. Brookmeyer R, Johnson E, Barry S. Modeling the incubation period of anthrax. *Stat Med.* 2005;24:531–542.
41. Walden J, Kaplan EH. Estimating time and size of bioterror attack. *Emerg Infect Dis.* 2004;10:1202–1205.

42. Barnett DJ, Balicer RD, Blodgett D, Fews AL, Parker CL, Links JM. The application of the Haddon matrix to public health readiness and response planning. *Environ Health Perspect.* 2005;113:561–566.
43. Meadows M. Project Bioshield: protecting Americans from terrorism. *FDA Consum.* 2004;38:32–33. Available at: http://www.fda.gov/fdac/features/2004/604_terror.html. Accessed August 15, 2006.
44. Quinn CP, Semenova VA, Elie CM, et al. Specific, sensitive, and quantitative enzyme-linked immunosorbent assay for human immunoglobulin G antibodies to anthrax toxin protective antigen. *Emerg Infect Dis.* 2002;8:1103–1110.
45. De BK, Bragg SL, Sanden GN, et al. A two-component direct fluorescent-antibody assay for rapid identification of *Bacillus anthracis*. *Emerg Infect Dis.* 2002;8:1060–1065.
46. Hoffmaster AR, Meyer RF, Bowen MD, et al. Evaluation and validation of a real-time polymerase chain reaction assay for rapid identification of *Bacillus anthracis*. *Emerg Infect Dis.* 2002;8:1178–1182.
47. Bell CA, UhlJR, Hadfield TL, et al. Detection of *Bacillus anthracis* DNA by LightCycler PCR. *J Clin Microbiol.* 2002;40:2897–2902.
48. Levine SM, Perez-Perez G, Olivares A, Yee H, Hanna BA, Blaser MJ. PCR-based detection of *Bacillus anthracis* in formalin-fixed tissue from a patient receiving ciprofloxacin. *J Clin Microbiol.* 2002;40:4360–4362.
49. Fasanella A, Losito S, Adone R, et al. PCR assay to detect *Bacillus anthracis* spores in heat-treated specimens. *J Clin Microbiol.* 2003;41:896–899.
50. Kaufmann AF. Observations on the occurrence of anthrax as related to soil type and rainfall. *Salisbury Med Bull Suppl.* 1990;68:16–17.
51. de Vos V. The ecology of anthrax in the Kruger National Park, South Africa. *Salisbury Med Bull Suppl.* 1990;68:19–23.
52. Wilson GS, Miles AA. *Topley and Wilson's Principles of Bacteriology and Immunity.* Vol 2. Baltimore, Md: Williams & Wilkins; 1955: 1940.
53. Davies JCA. A major epidemic of anthrax in Zimbabwe, III: distribution of cutaneous lesions. *Cent Afr J Med.* 1983;29:8–12.
54. Christopher GW, Agan MB, Cieslak TJ, Olson PE. History of US military contributions to the study of bacterial zoonoses. *Mil Med.* 2005;170(Suppl):39–48.
55. Glassman HN. World incidence of anthrax in man. *Public Health Rep.* 1958;73:22–24.
56. Fujikura T. Current occurrence of anthrax in man and animals. *Salisbury Med Bull Suppl.* 1990;68:1.
57. Shiferaw G. Anthrax in Wabessa village in the Dessie Zuria district of Ethiopia. *Rev Sci Tech.* 2004;23:951–956.
58. Maguina C, Flores del Pozo J, Terashima A, et al. Cutaneous anthrax in Lima, Peru: retrospective analysis of 71 cases, including four with a meningoencephalic complication. *Rev Inst Med Trop Sao Paulo.* 2005;47:25–30.
59. Leendertz FH, Ellerbrok H, Boesch C, et al. Anthrax kills wild chimpanzees in a tropical rainforest. *Nature.* 2004;430:451–452.
60. Liang X, Yu D. Identification of *Bacillus anthracis* strains in China. *J Appl Microbiol.* 1999;87:200–203.
61. Hugh-Jones M. 1996–97 Global anthrax report. *J Appl Microbiol.* 1999;87:189–191.
62. Brachman PS. Inhalation anthrax. *Ann N Y Acad Sci.* 1980;353:83–93.
63. Chang MH, Glynn MK, Groseclose SL. Endemic, notifiable bioterrorism-related diseases, United States, 1992–1999. *Emerg Infect Dis.* 2003;9:556–564.
64. El-kina AV. The epidemiology of a pulmonary form of anthrax. *Zh Mikrobiol Epidemiol Immunobiol.* 1971;48:112–116.

65. Plotkin SA, Brachman PS, Utell M, Bumford FH, Atchison MM. An epidemic of inhalation anthrax, the first in the twentieth century: I. clinical features. *Am J Med.* 1960;29:992–1001.
66. Davies JC. A major epidemic of anthrax in Zimbabwe. The experience at the Beatrice Road Infectious Diseases Hospital, Harare. *Cent Afr J Med (Zimbabwe)*. Part 1, 1982;28:291–298; Part 2, 1983;29:8–12; Part 3, 1985;31:176–180.
67. Koehler TM. *Bacillus anthracis* genetics and virulence gene regulation. *Curr Top Microbiol Immunol.* 2002;271:143–164.
68. Bail O. Quoted in: Sterne M. Anthrax. In: Stableforth AW, Galloway IA, eds. *Infectious Diseases of Animals*. Vol 1. London, England: Butterworths Scientific Publications; 1959: 22.
69. Ivins BE, Ezzell JW Jr, Jemski J, Hedlund KW, Ristroph JD, Leppla SH. Immunization studies with attenuated strains of *Bacillus anthracis*. *Infect Immun.* 1986;52:454–458.
70. Keppie J, Harris-Smith PW, Smith H. The chemical basis of the virulence of *Bacillus anthracis*, IX: its aggressins and their mode of action. *Brit J Exp Pathol.* 1963;44:446–453.
71. Drysdale M, Heninger S, Hutt J, Chen Y, Lyons CR, Koehler TM. Capsule synthesis by *Bacillus anthracis* is required for dissemination in murine inhalation anthrax. *Embo J.* 2005;24:221–227.
72. Schneerson R, Kubler-Kielb J, Liu TY, et al. Poly(gamma-D-glutamic acid) protein conjugates induce IgG antibodies in mice to the capsule of *Bacillus anthracis*: a potential addition to the anthrax vaccine. *Proc Natl Acad Sci U S A.* 2003;100:8945–8950.
73. Rhie GE, Roehrl MH, Mourez M, Collier RJ, Mekalanos JJ, Wang JY. A dually active anthrax vaccine that confers protection against both bacilli and toxins. *Proc Natl Acad Sci U S A.* 2003;100:10925–10930.
74. Chabot DJ, Scorpio A, Tobery SA, Little SF, Norris SL, Friedlander AM. Anthrax capsule vaccine protects against experimental infection. *Vaccine.* 2004;23:43–47.
75. Smith H, Keppie J. Observations on experimental anthrax: demonstration of a specific lethal factor produced in vivo by *Bacillus anthracis*. *Nature.* 1954;173:869–870.
76. Lincoln RE, Fish DC. Anthrax toxin. In: Montie TC, Kadis S, Ajl SJ, eds. *Microbial Toxins*. Vol 3. New York, NY: Academic Press; 1970: 361–414.
77. Stephen J. Anthrax toxin. In: Dorner F, Drews J, eds. *Pharmacology of Bacterial Toxins*. Oxford, England: Pergamon Press; 1986: 381–395.
78. Leppla SH. The anthrax toxin complex. In: Alouf JE, Freer JH, eds. *Sourcebook of Bacterial Protein Toxins*. London, England: Academic Press; 1991: 277–302.
79. Bartkus JM, Leppla SH. Transcriptional regulation of the protective antigen gene of *Bacillus anthracis*. *Infect Immun.* 1989;57:2295–2300.
80. Uchida I, Hornung JM, Thorne CB, Klimpel KR, Leppla SH. Cloning and characterization of a gene whose product is a trans-activator of anthrax toxin synthesis. *J Bacteriol.* 1993;175:5329–5338.
81. Sirard JC, Mock M, Fouet A. The three *Bacillus anthracis* toxin genes are coordinately regulated by bicarbonate and temperature. *J Bacteriol.* 1994;176:5188–5192.
82. Vietri NJ, Marrero R, Hoover TA, Welkos SL. Identification and characterization of a trans-activator involved in the regulation of encapsulation by *Bacillus anthracis*. *Gene.* 1995;152:1–9.
83. Bourgogne A, Drysdale M, Hilsenbeck SG, Peterson SN, Koehler TM. Global effects of virulence gene regulators in a *Bacillus anthracis* strain with both virulence plasmids. *Infect Immun.* 2003;71:2736–2743.

84. Mignot T, Mock M, Fouet A. A plasmid-encoded regulator couples the synthesis of toxins and surface structures in *Bacillus anthracis*. *Mol Microbiol*. 2003;47:917–927.
85. Hoffmaster AR, Ravel J, Rasko DA, et al. Identification of anthrax toxin genes in a *Bacillus cereus* associated with an illness resembling inhalation anthrax. *Proc Natl Acad Sci U S A*. 2004;101:8449–8454.
86. Avashia S, Riggins C, Lindley C, et al. Rapidly fatal *Bacillus cereus* pneumonia among metal workers—Texas 2003. Poster 398. Boston, Mass: 42nd Annual Meeting Infectious Disease Society of America, 2004.
87. Cone LA, Dreisbach L, Potts BE, Comess BE, Burleigh WA. Fatal *Bacillus cereus* endocarditis masquerading as an anthrax-like infection in a patient with acute lymphoblastic leukemia: case report. *J Heart Valve Dis*. 2005;14:37–39.
88. Mikesell P, Ivins BE, Ristroph JD, Dreier TM. Evidence for plasmid-mediated toxin production in *Bacillus anthracis*. *Infect Immun*. 1983;39:371–376.
89. Cataldi A, Labruyere E, Mock M. Construction and characterization of a protective antigen-deficient *Bacillus anthracis* strain. *Mol Microbiol*. 1990;4:1111–1117.
90. Pezard C, Berche P, Mock M. Contribution of individual toxin components to virulence of *Bacillus anthracis*. *Infect Immun*. 1991;59:3472–3477.
91. Kashiba S, Morishima T, Kato K, Shima M, Amano T. Leucotoxic substance produced by *Bacillus anthracis*. *Biken J*. 1959;2:97–104.
92. Stanley JL, Smith H. Purification of factor I and recognition of a third factor of the anthrax toxin. *J Gen Microbiol*. 1961;26:49–63.
93. Beall FA, Taylor MJ, Thorne CB. Rapid lethal effects in rats of a third component found upon fractionating the toxin of *Bacillus anthracis*. *J Bacteriol*. 1962;83:1274–1280.
94. O'Brien J, Friedlander A, Dreier T, Ezzell J, Leppla S. Effects of anthrax toxin components on human neutrophils. *Infect Immun*. 1985;47:306–310.
95. Wright GG, Read PW, Mandell GL. Lipopolysaccharide releases a priming substance from platelets that augments the oxidative response of polymorphonuclear neutrophils to chemotactic peptide. *J Infect Dis*. 1988;157:690–696.
96. Hoover DL, Friedlander AM, Rogers LC, Yoon IK, Warren RL, Cross AS. Anthrax edema toxin differentially regulates lipopolysaccharide-induced monocyte production of tumor necrosis factor alpha and interleukin-6 by increasing intracellular cyclic AMP. *Infect Immun*. 1994;62:4432–4439.
97. Tournier JN, Quesnel-Hellmann A, Mathieu J, et al. Anthrax edema toxin cooperates with lethal toxin to impair cytokine secretion during infection of dendritic cells. *J Immunol*. 2005;174:4934–4941.
98. Klimpel KR, Arora N, Leppla SH. Anthrax toxin lethal factor contains a zinc metalloprotease consensus sequence which is required for lethal toxin activity. *Mol Microbiol*. 1994;13:1093–1100.
99. Friedlander AM. Macrophages are sensitive to anthrax lethal toxin through an acid-dependent process. *J Biol Chem*. 1986;261:7123–7126.
100. Abrami L, Reig N, van der Goot FG. Anthrax toxin: the long and winding road that leads to the kill. *Trends Microbiol*. 2005;13:72–78.
101. Erwin JL, DaSilva LM, Bavari S, Little SF, Friedlander AM, Chanh TC. Macrophage-derived cell lines do not express proinflammatory cytokines after exposure to *Bacillus anthracis* lethal toxin. *Infect Immun*. 2001;69:1175–1177.
102. Pellizzari R, Guidi-Rontani C, Vitale G, Mock M, Montecucco C. Anthrax lethal factor cleaves MKK3 in macrophages and inhibits the LPS/IFN γ -induced release of NO and TNF α . *FEBS Lett*. 1999;462:199–204.

103. Agrawal A, Lingappa J, Leppla SH, et al. Impairment of dendritic cells and adaptive immunity by anthrax lethal toxin. *Nature*. 2003;424:329–334.
104. Warfel, JM, Steele AD, D'Agnillo F. Anthrax lethal toxin induces endothelial barrier dysfunction. *Am J Pathol*. 2005;166:1871–1881.
105. Fang H, Cordoba-Rodriguez R, Lankford CS, Frucht DM. Anthrax lethal toxin blocks MAPK kinase-dependent IL-2 production in CD4⁺ T cells. *J Immunol*. 2005;174:4966–4971.
106. Singh Y, Klimpel KR, Quinn CP, Chaudhary VK, Leppla SH. The carboxyl-terminal end of protective antigen is required for receptor binding and anthrax toxin activity. *J Biol Chem*. 1991;266:15493–15497.
107. Little SL, Lowe JR. Location of receptor-binding region of protective antigen from *Bacillus anthracis*. *Biochem Biophys Res Commun*. 1991;180:531–537.
108. Escuyer V, Collier RJ. Anthrax protective antigen interacts with a specific receptor on the surface of CHO-K1 cells. *Infect Immun*. 1991;59:3381–3386.
109. Bradley KA, Young JA. Anthrax toxin receptor proteins. *Biochem Pharmacol*. 2003;65:309–314.
110. Scobie HM, Young JA. Interactions between anthrax toxin receptors and protective antigen. *Curr Opin Microbiol*. 2005;8:106–112.
111. Bhatnagar R, Singh Y, Leppla SH, Friedlander AM. Calcium is required for the expression of anthrax lethal toxin activity in the macrophagelike cell line J774A.1. *Infect Immun*. 1989;57:2107–2114.
112. Klimpel KR, Molloy SS, Thomas G, Leppla SH. Anthrax toxin protective antigen is activated by a cell surface protease with sequence specificity and catalytic properties of furin. *Proc Natl Acad Sci U S A*. 1992;89:10277–10281.
113. Pimental RA, Christensen KA, Krantz BA, Collier RJ. Anthrax toxin complexes: heptameric protective antigen can bind lethal factor and edema factor simultaneously. *Biochem Biophys Res Commun*. 2004;322:258–262.
114. Abrami L, Liu S, Cosson P, Leppla SH, van der Goot FG. Anthrax toxin triggers endocytosis of its receptor via a lipid raft-mediated clathrin-dependent process. *J Cell Biol*. 2003;160:321–328.
115. Abrami L, Lindsay M, Parton RG, Leppla SH, van der Goot FG. Membrane insertion of anthrax protective antigen and cytoplasmic delivery of lethal factor occur at different stages of the endocytic pathway. *J Cell Biol*. 2004;166:645–651.
116. Ezzell JW Jr, Abshire TG. Serum protease cleavage of *Bacillus anthracis* protective antigen. *J Gen Microbiol*. 1992;138:543–549.
117. Panchal RG, Halverson KM, Ribot W, et al. Purified *Bacillus anthracis* lethal toxin complex formed in vitro and during infection exhibits functional and biological activity. *J Biol Chem*. 2005;280:10834–10839.
118. Ross JM. The pathogenesis of anthrax following the administration of spores by the respiratory route. *J Pathol Bacteriol*. 1957;73:485–494.
119. Dutz W, Kohout E. Anthrax. *Pathol Annu*. 1971;6:209–248.
120. Schwartz M. Cutaneous anthrax. *J Travel Med*. 2002;9:333.
121. Tutrone WD, Scheinfeld NS, Weinberg JM. Cutaneous anthrax: a concise review. *Cutis*. 2002;69:27–33.
122. Shieh WJ, Guarner J, Paddock C, et al. The critical role of pathology in the investigation of bioterrorism-related cutaneous anthrax. *Am J Pathol*. 2003;163:1901–1910.
123. Meade J. Images in emergency medicine. Cutaneous anthrax. *Ann Emerg Med*. 2004;43:664, 673.

124. Carucci JA, McGovern TW, Norton SA, et al. Cutaneous anthrax management algorithm. *J Am Acad Dermatol*. 2002;47:766–769.
125. Coban YK, Balik O, Boran C. Cutaneous anthrax of the hand and its reconstruction with a reverse-flow radial forearm flap. *Ann Plast Surg*. 2002;49:109–111.
126. Freedman A, Afonja O, Chang MW, et al. Cutaneous anthrax associated with microangiopathic hemolytic anemia and coagulopathy in a 7-month-old infant. *JAMA*. 2002;287:898–900.
127. Friedlander AM. Diagnosis and treatment of cutaneous anthrax. *JAMA*. 2002;288:43–44.
128. Khajehdehi P. Toxemic shock, hematuria, hypokalemia and hypoproteinemia in a case of cutaneous anthrax. *Mt Sinai J Med*. 2001;68:213–215.
129. Kaufmann AF, Dannenberg AL. Age as a risk factor for cutaneous human anthrax: evidence from Haiti, 1973–1974. *Emerg Infect Dis*. 2002;8:874–875. Available from: <http://www.cdc.gov/ncidod/EID/vol8no8/01-0207.htm>. Accessed August 18, 2006.
130. Franz DR, Jahrling PB, McClain DJ, et al. Clinical recognition and management of patients exposed to biological warfare agents. *Clin Lab Med*. 2001;21:435–473.
131. Inglesby TV, O’Toole T, Henderson DA, et al. Anthrax as a biological weapon, 2002: updated recommendations for management. *JAMA*. 2002;287:2236–2252.
132. Bartlett JG, Inglesby TV Jr, Borio L. Management of anthrax. *Clin Infect Dis*. 2002;35:851–858.
133. Spencer RC. *Bacillus anthracis*. *J Clin Pathol*. 2003;56:182–187.
134. Jernigan JA, Stephens DS, Ashford DA, et al. Bioterrorism-related inhalational anthrax: the first 10 cases reported in the United States. *Emerg Infect Dis*. 2001;7:933–944.
135. Centers for Disease Control and Prevention. Update: investigation of bioterrorism-related inhalational anthrax—Connecticut, 2001. *MMWR Morb Mortal Wkly Rep*. 2001;50:1049–1051.
136. Guarner J, Jernigan JA, Shieh WJ, et al. Pathology and pathogenesis of bioterrorism-related inhalational anthrax. *Am J Pathol*. 2003;163:701–709.
137. Hoffmaster AR, Meyer RF, Bowen MD, et al. Evaluation and validation of real-time polymerase chain reaction assay for rapid identification of *Bacillus anthracis*. *Emerg Infect Dis*. 2002;8:1178–1182.
138. Beatty ME, Ashford DA, Griffin PM, Tauxe RV, Sobel J. Gastrointestinal anthrax: review of the literature. *Arch Intern Med*. 2003;163:2527–2531.
139. Sejvar JJ, Tenover FC, Stephens DS. Management of anthrax meningitis. *Lancet Infect Dis*. 2005;5:287–295.
140. Krol CM, Uszynski M, Dillon EH, et al. Dynamic CT features of inhalational anthrax infection. *AJR Am J Roentgenol*. 2002;178:1063–1066.
141. Sirisanthana T, Brown AE. Anthrax of the gastrointestinal tract. *Emerg Infect Dis*. 2002; 8:649–651.
142. Kanafani ZA, Ghossain A, Sharara AI, Hatem JM, Kanj SS. Endemic gastrointestinal anthrax in 1960s Lebanon: clinical manifestations and surgical findings. *Emerg Infect Dis*. 2003;9:520–525.
143. Turnbull PCB, Leppla SH, Broster MG, Quinn CP, Melling J. Antibodies to anthrax toxin in humans and guinea pigs and their relevance to protective immunity. *Med Microbiol Immunol*. 1988;177:293–303.
144. Buchanan TM, Feeley JC, Hayes PS, Brachman PS. Anthrax indirect microhemagglutination test. *J Immunol*. 1971;107:1631–1636.

145. Sirisanthana T, Nelson KE, Ezzell J, Abshire TG. Serological studies of patients with cutaneous and oral-oro-pharyngeal anthrax from northern Thailand. *Am J Trop Med Hyg.* 1988;39:575–581.
146. Harrison LH, Ezzell JW, Abshire TG, Kidd S, Kaufmann AF. Evaluation of serologic tests for diagnosis of anthrax after an outbreak of cutaneous anthrax in Paraguay. *J Infect Dis.* 1989;160:706–710.
147. Shlyakhov EN, Rubinstein E. Human live anthrax vaccine in the former USSR. *Vaccine.* 1994;12:727–730.
148. Pfisterer RM. Retrospective verification of the diagnosis of anthrax by means of the intracutaneous skin test with the Russian allergen anthraxin in a recent epidemic in Switzerland. *Salisbury Med Bull Suppl.* 1990;68:80.
149. Centers for Disease Control and Prevention. Update: investigation of bioterrorism-related anthrax and interim guidelines for exposure management and antimicrobial therapy. *MMWR Morb Mortal Wkly Rep.* 2001;50:909–919.
150. Shepard CW, Soriano-Gabarro M, Zell ER, et al. Antimicrobial postexposure prophylaxis for anthrax: adverse events and adherence. *Emerg Infect Dis.* 2002;8:1124–1132.
151. Jefferds MD, Laserson K, Fry AM, et al. Adherence to antimicrobial inhalational anthrax prophylaxis among postal workers, Washington, DC, 2001. *Emerg Infect Dis.* 2002;8:1138–1144.
152. Navas E. Problems associated with potential massive use of antimicrobial agents as prophylaxis or therapy of a bioterrorist attack. *Clin Microbiol Infect.* 2002;8:534–539.
153. Williams JL, Noviello SS, Griffith KS, et al. Anthrax postexposure prophylaxis in postal workers, Connecticut, 2001. *Emerg Infect Dis.* 2002;8:1133–1137.
154. Blank S, Moskin LC, Zucker JR. An ounce of prevention is a ton of work: mass antibiotic prophylaxis for anthrax, New York City, 2001. *Emerg Infect Dis.* 2003;9:615–622.
155. Friedlander AM, Heine H. *Bacillus anthracis*. In: Yu VL, Weber R, Raoult D, eds. *Antimicrobial Therapy and Vaccines*. New York, NY: Apple Trees Productions, LLC; 2002.
156. Bryskier A. *Bacillus anthracis* and antibacterial agents. *Clin Microbiol Infect.* 2002;8:467–478.
157. Torres-Tortosa M, Caballero-Granado FJ, Moreno I, Canueto J. Antimicrobial therapy for anthrax. *Eur J Clin Microbiol Infect Dis.* 2002;21:696.
158. Tasyaran MA, Erol S. Penicillin therapy for anthrax: a reply to Torres-Tortosa et al. *Eur J Clin Microbiol Infect Dis.* 2002;21:697.
159. Beharry Z, Chen H, Gadachanda VR, Buynak JD, Palzkill T. Evaluation of penicillin-based inhibitors of the class A and B β -lactamases from *Bacillus anthracis*. *Biochem Biophys Res Commun.* 2004;313:541–545.
160. Chen Y, Tenover FC, Koehler TM. β -Lactamase gene expression in a penicillin-resistant *Bacillus anthracis* strain. *Antimicrob Agents Chemother.* 2004;48:4873–4877.
161. Gold H. Anthrax: a report of one hundred seventeen cases. *AMA Arch Intern Med.* 1955;96:387–396.
162. Lightfoot NF, Scott RJD, Turnbull PCB. Antimicrobial susceptibility of *Bacillus anthracis*. *Salisbury Med Bull Suppl.* 1990;68:95–98.
163. Doganay M, Aydin N. Antimicrobial susceptibility of *Bacillus anthracis*. *Scand J Infect Dis.* 1991;23:333–335.
164. Mikesell P. Major, Medical Service, US Army. Investigator, Division of Bacteriology, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, Md. Personal communication, January 1991.
165. Patt HA, Feigin RD. Diagnosis and management of suspected cases of bioterrorism; a pediatric perspective. *Pediatrics.* 2002;109:685–692.

166. Friedlander AM, Welkos SL, Pitt MLM, et al. Postexposure prophylaxis against experimental inhalation anthrax. *J Infect Dis.* 1993;167:1239–1243.
167. Brookmeyer R, Johnson E, Bollinger R. Public health vaccination policies for containing an anthrax outbreak. *Nature.* 2004;432:901–904.
168. Fowler RA, Sanders GD, Bravata DM, et al. Cost-effectiveness of defending against bioterrorism: a comparison of vaccination and antibiotic prophylaxis against anthrax. *Ann Intern Med.* 2005;142:601–610.
169. Vietri NJ, Purcell BK, Lawler JV, et al. Short-course postexposure antibiotic prophylaxis combined with vaccination protects against experimental inhalational anthrax. *Proc Natl Acad Sci U S A.* 2006;103:7813–7816.
170. Ivins BE, Welkos SL. Recent advances in the development of an improved, human anthrax vaccine. *Eur J Epidemiol.* 1988;4:12–19.
171. Little SF, Knudson GB. Comparative efficacy of *Bacillus anthracis* live spore vaccine and protective antigen vaccine against anthrax in the guinea pig. *Infect Immun.* 1986;52:509–512.
172. Turnbull PC, Broster MG, Carman JA, Manchee RJ, Melling J. Development of antibodies to protective antigen and lethal factor components of anthrax toxin in humans and guinea pigs and their relevance to protective immunity. *Infect Immun.* 1986;52:356–363.
173. Ivins BE, Welkos SL. Cloning and expression of the *Bacillus anthracis* protective antigen gene in *Bacillus subtilis*. *Infect Immun.* 1986;54:537–542.
174. Little SF, Webster WM, Ivins BE, Fellows PF, Norris SL, Andrews GP. Development of an in vitro-based potency assay for anthrax vaccine. *Vaccine.* 2004;22:2843–2852.
175. Centers for Disease Control and Prevention. Use of anthrax vaccine in the United States. Recommendations of the Advisory Committee on Immunization Practices. *MMWR Recomm Rep.* 2000;49:1–20.
176. Centers for Disease Control and Prevention. Use of anthrax vaccine in response to terrorism: supplemental recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep.* 2002;51:1024–1026.
177. Sever JL, Brenner AI, Gale AD, et al. Safety of anthrax vaccine: an expanded review and evaluation of adverse events reported to the Vaccine Adverse Event Reporting System (VAERS). *Pharmacoepidemiol Drug Saf.* 2004;13:825–840.
178. Friedlander AM, Pittman PR, Parker GW. Anthrax vaccine: evidence for safety and efficacy against inhalational anthrax. *JAMA.* 1999;282:2104–2106.
179. Pittman PR, Gibbs PH, Cannon TL, Friedlander AM. Anthrax vaccine: short-term safety experience in humans. *Vaccine.* 2001;20:972–978.
180. Pittman PR, Kim-Ahn G, Pifat DY, et al. Anthrax vaccine: immunogenicity and safety of a dose-reduction, route-change comparison study in humans. *Vaccine.* 2002;20:1412–1420.
181. Pittman PR. Aluminum-containing vaccine associated adverse events: role of route of administration and gender. *Vaccine.* 2002;20(Suppl 3):S48–S50.
182. Lange JL, Lesikar SE, Rubertone MV, Brundage JF. Comprehensive systematic surveillance for adverse effects of anthrax vaccine adsorbed, US Armed Forces, 1998–2000. *Vaccine.* 2003;21:1620–1628.
183. Wasserman GM, Grabenstein JD, Pittman PR, et al. Analysis of adverse events after anthrax immunization in US Army medical personnel. *J Occup Environ Med.* 2003;45:222–233.
184. Tierney BC, Martin SW, Franzke LH, et al. Serious adverse events among participants in the Centers for Disease Control and Prevention’s Anthrax Vaccine and Antimicrobial Availability Program for persons at risk for bioterrorism-related inhalational anthrax. *Clin Infect Dis.* 2003;37:905–911.

185. Grabenstein JD. Anthrax vaccine: a review. *Immunol Allergy Clin North Am.* 2003;23:713–730.
186. Martin SW, Tierney BC, Aranas A, et al. An overview of adverse events reported by participants in CDC's anthrax vaccine and antimicrobial availability program. *Pharmacoepidemiol Drug Saf.* 2005;14:393–401.
187. Friedlander AM, Welkos SL, Ivins BE. Anthrax vaccines. *Curr Top Microbiol Immunol.* 2002;271:33–60.
188. Johnson-Winegar A. Comparison of enzyme-linked immunosorbent and indirect hemagglutination assays for determining anthrax antibodies. *J Clin Microbiol.* 1984;20:357–361.
189. Pittman PR. Lieutenant Colonel, Medical Corps, US Army. Chief, Clinical Investigation, Medical Division, US Army Medical Research Institute of Infectious Diseases, Fort Detrick, Md. Personal communication, January 1994.
190. Hambleton P, Carman JA, Melling J. Anthrax: the disease in relation to vaccines. *Vaccine.* 1984;2:125–132.
191. Brachman PS, Gold H, Plotkin SA, Fekety FR, Werrin M, Ingraham NR. Field evaluation of a human anthrax vaccine. *Am J Public Health.* 1962;52:632–645.
192. Wright GG, Green TW, Kanode RG Jr. Studies on immunity in anthrax. V. Immunizing activity of alum-precipitated protective antigen. *J Immunol.* 1954;73:387–391.
193. Ivins BE, Welkos SL, Little SF, Crumrine MH, Nelson GO. Immunization against anthrax with *Bacillus anthracis* protective antigen combined with adjuvants. *Infect Immun.* 1992;60:662–668.
194. Ivins BE, Fellows PF, Nelson GO. Efficacy of a standard human anthrax vaccine against *Bacillus anthracis* spore challenge in guinea-pigs. *Vaccine.* 1994;12:872–874.
195. Ivins BE, Fellows PF, Pitt MLM, et al. Efficacy of a standard human anthrax vaccine against *Bacillus anthracis* aerosol spore challenge in rhesus monkeys. *Salisbury Med Bull.* 1996;87:125–126.
196. Ivins BE, Pitt ML, Fellows PF, et al. Comparative efficacy of experimental anthrax vaccine candidates against inhalation anthrax in rhesus macaques. *Vaccine.* 1998;16:1141–1148.
197. Pitt ML, Little S, Ivins BE, et al. In vitro correlate of immunity in an animal model of inhalational anthrax. *J Appl Microbiol.* 1999;87:304.
198. Fellows PF, Linscott MK, Ivins BE, et al. Efficacy of a human anthrax vaccine in guinea pigs, rabbits, and rhesus macaques against challenge by *Bacillus anthracis* isolates of diverse geographical origin. *Vaccine.* 2001;19:3241–3247.
199. Puziss M, Wright GG. Studies on immunity in anthrax. X. Gel-adsorbed protective antigen for immunization of man. *J Bacteriol.* 1963;85:230–236.
200. Sever JL, Brenner AI, Gale AD, et al. Safety of anthrax vaccine: a review by the Anthrax Vaccine Expert Committee (AVEC) of adverse events reported to the Vaccine Adverse Event Reporting System (VAERS). *Pharmacoepidemiol Drug Saf.* 2002;11:189–202.